

The Voice

The Voice

VOICE OF THE EDITOR

Mary Sandage, PhD

IN THIS ISSUE:

THE VOICE OF
THE EDITOR 1

BY EDITOR
MARY SANDAGE,
PHD

WORKING WITH 2-4
INJURED
PEDIATRIC
PERFORMERS

BY MAIA
BRADEN, MS,
CCC-SLP

WHAT DO WE 5-8
KNOW OF VOCAL
FOLD VIBRATORY
NOTION
IN CHILDREN?

BY RITA R.
PATEL, PHD,
CCC-SLP

2016 8
SYMPOSIUM
MASTER CLASS
LINK & PHOTOS

PEDIATRIC 9-10
VOICE THERAPY:
HOW CHILDREN
LEARN BEST

BY LESLIE S.
KESSLER, MA,
CCC-SLP

2016 PHOTOS 11-14
SYMPOSIUM
POSTER SESSION

STROBOSCOPY
OSCARS

PRESENTATION
OUTREACH

AWARD
PRESENTATIONS

CONTACT

15

Assessing, Treating and Training VOICES OF CHILDREN

BY MARY SANDAGE, PHD

It seems that most of our attention in the world of voice habilitation and rehabilitation focuses on the fully developed mature voices of our adult clients and voice students. For many of us in practice, we also encounter young children and adolescents with voice impairment, both congenital and acquired.

Assessing, treating and training the voices of children and adolescents requires special consideration. In this pediatric-focused issue I have invited researchers and practitioners who specialize in pediatric voice to share their expertise. Rita Patel, PhD, provides an update on recent high speed laryngeal imaging findings in pre-adolescent

children. As it turns out the old adage that children are not little adults appears to be true for vocal fold vibratory characteristics.

Maia Braden and Leslie Kessler both provide important clinical perspectives on child learning. Ms. Kessler considers learning differences in young children and delineates important adaptations to make in resonant voice treatment approaches. Ms. Braden skillfully describes the moving target that is the adolescent performing voice.



VOICE OF THE EDITOR
MARY SANDAGE, PHD

*Assistant Professor
Dept. of Communication Disorders
Auburn University*



*Voice
of the Editor*

WORKING WITH INJURED PEDIATRIC PERFORMERS

BY MAIA BRADEN, MS,
CCC-SLP

Like adult performers, pre-teen and teenage singers are vocal athletes and face similar risks for injury. However, young performers also present a separate set of challenges from their adult counterparts. They do not have fully mature instruments to begin with, are developing mature vocal technique, and learn differently from adults. Their vocal demands are different, and they have different barriers and supports to therapy than adults.

LARYNGEAL DEVELOPMENT AND VOICE MATURATION

The adolescent voice is a moving target for a variety of reasons. The mature 5-layer structure of vocal folds is not fully differentiated until the teen years, between 14 and 17 years of age, and the length of the membranous vocal fold in

relation to the cartilaginous portion increases with age.¹ Male pubertal changes are marked by rapid growth in vocal fold length and in vocal tract length, and females undergo voice change as well, with a lowering and then expansion of range.²⁻⁴ If we consider how singing involves finely coordinated adjustments of vocal fold length and tension, vocal tract dimensions, respiratory control, and use of articulators, rapid changes in any of these, as well as developing fine motor control are likely to lead to instability. It is normal in adolescence to experience change in voice type, periods of instability, and periods of increased difficulty with register transitions, due to these changes occurring.⁵ Sometimes attempts to navigate these changes can result in maladaptive compensations predisposing singers to vocal injuries.

TRAINING AND TECHNIQUE

When we see adult injured singers, most have established singing technique. Conversely, most very young singers are still in the infancy of developing their technical skills. They may have a vague idea of optimal breathing for singing, alignment, resonance, and coordination of the subsystems of respiration, phonation and resonance. More often, they are going entirely on instinct. Developing singers often try to imitate much older performers, and may receive a great deal of positive reinforcement for doing this – it seems as if listeners are always inordinately impressed when they hear a 10-year-old who sounds like a 30-year-old. Young singers (in fact all singers) should be encouraged to sound like the best version of themselves, and both singing teachers and speech pathologists who

(Continued on page 3)

The Voice of the
Singer / Pediatric Therapist

WORKING WITH INJURED PEDIATRIC PERFORMERS, CONTINUED

(Continued from page 2)

work with singers should address the basics of healthy vocal technique.

LEARNING

Pre-teens and teens are in a period of very rapid learning, and they are accustomed to learning new things every day. In this way, they can be easier to work with than adults. However, we need to keep in mind some of the learning differences in this age group. Their pre-frontal cortex is still developing, and while executive function is improving, they often struggle with impulse control and prioritization.⁶ Adolescents learn best when the learning is meaningful and authentic, and when they have opportunities to set their own goals and monitor their own progress. This means asking them what they hope to accomplish, and asking them to judge their own performance and adherence to therapy tasks. Learning needs to be presented in multiple ways (written, modeled, experienced, recorded) to support their best learning.

VOCAL DEMANDS

Teens often have extremely full schedules both at school and after. They may be playing a sport, performing in a play, working a part-

time job, and singing in a band, at church, or in a choir. Combined with the lower impulse control that comes with adolescence and the natural desire to be social and talk, vocal demands of adolescents can be extremely high. Therapy should focus on healthy production in *all* voice use. After all, we don't put our "singing" voice away and take out our "speaking" voice. In fact, teenagers may be using a healthier production technique when singing than when speaking. Given all of their competing vocal demands, we discuss voice prioritization rather than limitation, and help them to see their voice as a long-term project. When being told what to do and what not to do, they are likely to respond defensively, but

"The adolescent voice is a moving target for a variety of reasons."

when approached as collaborative problem-solving, they are more likely to make healthy choices on their own.

EXTERNAL SUPPORTS

While parents may be a primary support in therapy, peers and other adults also play a supporting role. Other adult supports include choir directors, teachers, voice teachers and coaches, and theater directors. Collaboration with everyone involved in a patient's voice care is necessary to



MAIA BRADEN, MS,
CCC-SLP

*Singer
Pediatric Voice Manager
UW Madison Voice and
Swallow Clinics
Madison, WI*

make sure that the patient is getting all they need. Conversations about healthy vocal technique, vocal expectations, and making sure that the same

recommendations are being reinforced in all settings are vital.

CONCLUSION

In working with pre-teen and teenage performers, we have the unique opportunity to set them up for success in the future. We still have much to learn about how the voice develops, and as we do we can serve this population better. With these factors in mind, we should not shy away from working with this popu-

(Continued on page 4)

WORKING WITH INJURED PEDIATRIC PERFORMERS, CONTINUED

(Continued from page 3)

lation, but should use our knowledge of vocal rehabilitation to help them recover and develop into successful adult performers.



1. Hirano M, Kurita S, Nakashima T. Growth, development and aging of human vocal folds. In: Bless D, Abbs J, eds. *Vocal Fold Physiology*. San Diego, CA: College-Hill Press; 1983.
2. Willis EC, Kenny DT. Relationship between weight, speaking fundamental frequency, and the appearance of phonational gaps in the adolescent male changing voice. *J Voice*. 2008;22(4):451-471.
3. Sapienza CM, Ruddy BH, Baker S. Laryngeal structure and function in the pediatric larynx: clinical applications. *Lang Speech Hear Serv Sch*. 2004;35(4):299-307.
4. Gackle L. Understanding voice transformation in female adolescents. In: Thurman L, Welch G, eds. *Bodymind and Voice: Foundations of voice education*. Collegeville: The Voice Care Network; 2000:739-744.
5. Boltezar IH, Burger ZR, Zargi M. Instability of voice in adolescence: pathologic condition or normal developmental variation? *J Pediatr*. 1997;130(2):185-190.
6. Blakemore S-J, Choudhury S. Development of the adolescent brain: implications for executive function and social cognition. *Journal of Child Psychology and Psychiatry*. 2006;47(3-4):296-312.

Maia Braden, MS, CCC-SLP is the pediatric voice manager at the UW Madison Voice and Swallow Clinics in Madison, WI. She holds a BA in vocal performance and an MS in communicative disorders from the University of Wisconsin, Madison. She presents extensively on voice and voice disorders locally and nationally. She has published in peer-reviewed journals on topics related to voice disorders, and has a special interest in the pediatric performing voice. She has performed in choral, musical theater, folk, and opera settings in a group and as a soloist.

CALL FOR PAPERS

46TH ANNUAL SYMPOSIUM:

CARE OF THE PROFESSIONAL VOICE

Wednesday, May 31 through Sunday, June 4, 2017

The Westin Philadelphia Hotel

The Voice Foundation is seeking abstracts for oral and poster presentations as well as proposals for workshops.

Abstract submission is available online at [Abstract Submission](#)

Opening: September 1, 2016

Abstract Submission Workshops Deadline: October 15, 2016

Abstract Submission Papers/Posters Deadline: October 31, 2016

Notification of Acceptance: January 31, 2017

WHAT DO WE KNOW OF VOCAL FOLD VIBRATORY MOTION IN CHILDREN?

BY RITA R. PATEL, PHD, CCC-SLP

Voice disorders occur across the lifespan, are disabling and compromise quality of life. Conservative estimates of children suffering from voice disorders suggest that over 1 million or 6-9% children experience voice problems nationwide.¹ Other estimates have placed the occurrence of voice disorders in school-aged children as high as 16%.² Because a voice disorder results in an alteration of voice quality (i.e. dysphonia or hoarseness), an untreated voice disorder can substantially hinder children's academic performance and overall psychological well-being.³⁻⁵ Hence early identification and treatment of voice disorders in children is essential. The clinical evaluation and treatment of voice disorders in children requires a clinician to assess the vocal fold structure and function. Children not only have smaller laryngeal structures but also differ substantially in terms of the vocal fold layered structures.⁶⁻⁹ The five layers of the vocal folds are not fully developed in children, but mature only after puberty. Since structurally, the children's vocal folds are not simply a smaller-

sized version of the adult vocal folds, information on vibratory motion from adults cannot be directly used for interpretation of children's vocal fold vibratory motion.

A comprehensive evaluation of vocal fold function involves the evaluation of the pliability and the cycle-to-cycle motion of the vocal folds over a period of time. As the vocal fold vibrations are too rapid for the human eye to detect, it is critical to slow the motion of the vocal folds in order to evaluate their function. In children, the vocal folds vibrate anywhere between 218 to 329 times per second.¹⁰ Such rapid vibrations are difficult for the human eye to perceive with simple endoscopy, which uses a straight halogen light. The current state-of-the-art clinical assessment technique is stroboscopy: a flashing light and an endoscope are used to observe *apparent* cycles of vocal fold vibrations.¹⁰ The vocal fold vibratory cycle consists of three phases: the opening phase, the closing phase, and the closed phase (Figure 1). Stroboscopy allows us to view the vocal fold motion at a slower rate by illuminating only parts of succes-



RITA R. PATEL, PHD,
CCC-SLP


*Assistant professor
Department of Speech
and Hearing Sciences
Indiana University*

sive cycles—but at the price of substantial information loss. That is, stroboscopy does not provide the viewer with information from each cycle of vocal fold vibration, but instead illuminates only parts of successive cycles. Moreover, it does not work well for severely impaired voice qualities¹¹ or short phonations of 2-3 seconds,¹² both of which are frequently encountered in clinical examinations in children. A rela-

The Voice of the Therapist

(Continued on page 6)

Figure 1: One glottal cycle in a 10 year old female child



(Continued from page 5)

tively new method of laryngeal high-speed videoendoscopy can image the vocal folds at 8000 frames per second: an enormous advance on stroboscopy, which images the vocal folds at 30 frames per second.¹³ Because of this increased temporal resolution, high-speed videoendoscopy allows for effective evaluation of the entire range of voice qualities and durations encountered in the pediatric population.

Our findings from both qualitative and quantitative evaluation of cycles of vocal fold vibrations have revealed that vibratory motion in children is not a small version of adult oscillation, but is functionally different. Our studies have focused on evaluation of steady-state sustained phonation at typical pitch and loudness in pre-pubertal children between the ages of 5 and 11 years. We used high-speed videoendoscopy to image the vocal folds, allowing us to view individual cycles of vocal fold vibrations and then slow them down for further viewing

and analysis. We found that children's vocal fold vibrations differ from those of adults in several ways.

First, children have a higher prevalence of a posterior phonatory gap (girls = 85%, boys = 68%) compared to adults (females = 75%, males = 54%).¹⁴ The posterior gap when present in children also appears to be more like a 'diamond' shape, i.e. extending not only in the posterior direction, but also in the anterior direction from the vocal process. This means that the presence of a posterior gap is not an indication of an abnormality, but part of the normal development of vibratory motion in children.

Second, children's vocal fold vibratory motion is characterized by a longer opening phase and a relatively shorter duration of the closing phase compared to vocally healthy young adult male and females.^{15,16} Typically developing children also showed a significantly larger amplitude-to-length ratio compared to adult females.¹⁵ These findings suggest that vocal fold

movements of typically developing children are more similar to those of adult males than adult females. Further, the excessively large amplitude-to-length ratio and the relatively shorter closing phase interval may explain why children have a harder time varying pitch and loudness independently of each other.

Interestingly, children also have a longer longitudinal phase delay between the anterior and the posterior parts of the vocal fold during the opening phase. The closing phase in children is more shutter-like without the longitudinal phase delay; more than adults.¹⁶ These differences suggest that children might have a different mechanism for opening and closing the vocal folds than adults do.

Third, children's vocal fold vibrations are more inconsistent from one cycle to another, i.e. children exhibit greater cycle-to-cycle variability. More specifically, children exhibited greater variability in both amplitude and time periodicity compared to adult men; however, they

(Continued on page 7)

(Continued from page 6)

were similar in terms of these parameters to adult females.¹⁷ Furthermore, the use of the Phonovibrogram image-based analysis technique to investigate movement of the anterior and posterior glottis during the opening and closing phases of the glottal cycle revealed greater differences in cycle-to-cycle variability along the anterior glottis during the closing phase. This increased variability suggests that the closing phase is more variable compared to the opening phase.¹⁶ The presences of these aperiodicities in vibratory motion would need to be taken into account when interpreting vocal fold vibratory motion during a clinical evaluation.

In conclusion, these findings indicate that vocal fold motion in children is complex and not easily predicted from adult normative data. We need to develop child-specific norms, because adult-like vibratory motion is not completely established by 11 years of age but continues to develop during this period.



References:

1. Wilson DK. Voice problems in children. 3rd. ed. Baltimore/London: Williams and Wilkins; 1987.
2. Kallvik E, Lindstrom E,

Dr. Rita Patel is an assistant professor at the Department of Speech and Hearing Sciences, Indiana University. Her research and clinical expertise focuses on addressing the vibratory kinematics of normal and abnormal voice in the adult and pediatric population, with high-speed videoendoscopy. Prior to starting her position at Indiana University, she was an Assistant Professor and Director Voice Clinic at the University of Kentucky for four years. Since her doctoral degree in 2006 Dr. Patel has been awarded 7 extramural grants from National agencies including ASHA and NIH; and 4 intramural grants. Dr. Patel has received 15 awards and 2 fellowships for her work in the area of voice. Currently she is the Associate Editor for the journal *Language, Speech, and Hearing Services in Schools*. She is also the coordinator of ASHAs Special Interest Group 3, Voice and Voice Disorders.

Holmqvist S, Lindman J, Simberg S. Prevalence of Hoarseness in School-aged Children. *J Voice* 2014;29:260.e1-.e19.

3. Connor NP, Cohen SB, Theis SM, Thibeault SL, Heatley DG, Bless DM. Attitudes of children with dysphonia. *J Voice* 2008;22:197-209.

4. Ruddy BH, Sapienza CM. Treating voice disorders in the school-based setting: working within the framework of IDEA. *Language, Speech, and Hearing Services in Schools* 2004;35:327-32.

5. Lass NJ, Ruscello DM, Bradshaw KH, Blankenship BL. Adolescents' perceptions of normal and voice-disordered children. *J Commun Disord* 1991;24:267-74.

6. Kahane JC. Growth of the human prepubertal and pubertal larynx. *J Speech Hear Res* 1982;25:446-55.

7. Hirano M, Kurita S, Nakashima T. Growth, Development, and Aging of Human Vocal

Fold: College-Hill Press, San Diego, California; 1983.

8. Hartnick CJ, Rehbar R, Prasad V. Development and maturation of the pediatric human vocal fold lamina propria. *Laryngoscope* 2005;115:4-15.

9. Sato K, Hirano M, Nakashima T. Fine structure of the human newborn and infant vocal fold mucosae. *Ann Otol Rhinol Laryngol* 2001;110:417-24.

10. Bless D, Glaze LE, Lowery-Biever D, Campos G, Peppard RC. Stroboscopic, acoustic, aerodynamic, and perceptual analysis of voice production in normal speaking adults. *NCVS Status and Progress Report* 1993;4:121-34.

11. Patel, Dailey S, Bless D. Comparison of high-speed digital imaging with stroboscopy for laryngeal imaging of glottal disorders. *Ann Otol Rhinol Laryngol* 2008;117:413-24.

12. Tigges M, Wittenberg T,

(Continued on page 8)

(Continued from page 7)

Mergell P, Eysholdt U. Imaging of vocal fold vibration by digital multi-plane kymography. *Comput Med Imaging Graph* 1999;23:323-30.

13. Deliyski DD, Hillman RE. State of the art laryngeal imaging: research and clinical implications. *Curr Opin Otolaryngol Head Neck Surg* 2010;18:147-52.

14. Patel, Dixon A, Richmond A, Donohue KD. Pediatric high

speed digital imaging of vocal fold vibration: A normative pilot study of glottal closure and phase closure characteristics. *Int J Pediatr Otorhinolaryngol* 2012;76:954-9.

15. Patel R, Donohue KD, Unnikrishnan H, Kryscio RJ. Kinematic measurements of the vocal-fold displacement waveform in typical children and adult populations: quantification of high-speed endoscopic videos. *J Speech Lang Hear Res* 2015;58:227-40.

16. Döllinger M, Dubrovskiy D, Patel R. Spatiotemporal analysis of vocal fold vibrations between children and adults. *Laryngoscope* 2012;122:2511-8.

17. Patel, Dubrovskiy D, Dollinger M. Characterizing vibratory kinematics in children and adults with high-speed digital imaging. *J Speech Lang Hear Res* 2014;57:S674-S86.

Link to [Symposium Master Class](#) - The Return of Dolora Zajick!



June 4th, 2016 Master Class Singers-MDs:
Isabel Garcia-Lopez, Josef Schlömicher-Thier, Jennifer Rodney

PEDIATRIC VOICE THERAPY: HOW CHILDREN LEARN BEST

BY LESLIE S. KESSLER, MA,
CCC-SLP

When I started providing pediatric voice therapy I knew it was necessary to develop a true resonant voice therapy program which mimicked the fine resonant voice therapy we provide our adult voice patients.

As a well-trained clinician I immediately began to facilitate awareness of vibrations in the front of their little faces. I did not use the word "vibrations" because I thought children would not understand this big word. I began using the word "buzzy" like a bee.

I felt so smart and inventive because my little ones so quickly reported feeling the "buzzy" on their lips, nose in their mouths and even their cheeks. Fortunately, I became skeptical of this immediate learning because adults, who have knowledge and experience with vibrations, did not catch on this quickly.

One day I started asking "do you feel the buzzy on your elbow?" "oh yes" "and on your knee?" "yes my knee is very buzzy!"

Lesson learned...most children have never had enough experience or any experience with cognitive awareness of vibrations and so cannot be

expected to determine presence or lack of presence of vibrations.

Proprioception of vibratory sensations is the foundation of resonant voice therapy. We needed a pediatric voice therapy which first addressed the child's lack of specific prior sensory, cognitive and language-based knowledge and experience, already present in the adult.

But, to facilitate this specific knowledge in children it is important to acknowledge that children learn differently from adults, and techniques that are detrimental to motor learning in adults (imagery and metaphors, for example) are key to child learning.

There is research specifically targeting child skill acquisition and cognitive/motor learning which highlights some differences in child and adult learning.

USE OF METAPHORS - ADULT POPULATION

Katherine Verdolini-Abbot PhD, CCC-SLP et al has published robust findings reporting that use of metaphors in teaching adults inhibits motor learning.

CHILD POPULATION

Laura S. DeThorne et al, (2009) reveals the results of their study indicated 1) greater accuracy and less variability with external focus for

tongue and hand control tasks, consistent with previous limb studies in adults and 2) pairing of specific speech sounds with meaningful environmental sounds, objects or events may lessen the load on the child's working memory during complex motor tasks. This metaphoric language engages visual processing skills or episodic memory and unburdens reliance on phonological processing. Examples of these metaphors are /u/ for monkey, /m/ for the yummy sound. I chose "buzzy bee" and "croaky frog" for Buzzy Child.

Sawada, MOM, & Ishii (2002) reported that 4 - 5:11 year old children were unable to attend to particular and relevant features of modeled actions, to learn a sequence of five dance steps and performed better with a verbal model. The verbal model chosen were specific metaphors. The rationale, according to Weiss, Ebbeck, & Wiese-Bjornstal (1993) "it is critical for young children to understand a physical task by associating it with a label and then relate the label to their own experiences." Therefore, metaphoric language is recommended when teaching motor skills. Results of this study indicated that metaphorical verbal instruction was more effective than demonstration to support



LESLIE S. KESSLER,
MA, CCC-SLP

*Owner and Director
Language & Voice
Experience Clinics
Washington, D.C.*

The
Voice
of
the
Clinician

(Continued on page 10)

ACOUSTIC PEDAGOGY, CONTINUED

(Continued from page 9)

young children's motor learning of sequential dance steps skills (4-5:11). Older children (7-8:11) benefited from both forms of instruction. According to these findings specific metaphorical instruction is appropriate to use with 4 - 8:11 year old children as an effective teaching and "learning strategy."

Research results pertaining to children, strongly suggest that the use of metaphorical instruction benefits young children where adults do not benefit from metaphorical instruction and perform more poorly. There seems to be no clear-cut age benchmark identifying "young children" but research suggests children three – seven chronological/developmental years are the children who



reap the most benefit from metaphorical instruction, although some studies report these same benefits through 9 or 10 years. Based on these and other cognitive-developmental factors in motor learning and modeling, specific metaphorical verbal instruction as described above or verbal rehearsal apparently influences modeling of a movement sequence in young

children, as it enables young children to attend to task-relevant information (Sawada et al, 2002) thus affording them effective strategies for motor learning. Somewhere between 12-17 years old there appears to be a shift from the child model of learning, to adult the model of learning.

SKILL ACQUISITION IN CHILDREN - HOW LONG DOES IT TAKE?

Jacoby et al. (2002) conducted a study to determine the average number of treatment units (15 minutes per unit) needed to achieve improvements in communication (voice being part of communication). The subjects in this study were 234 children between the ages 3-6 years old. The authors used a 7 level functional rating scale derived from the NOMS. (Level 7 was the discharge level.) The overall findings were

that children required, on average, 20 hours of therapy to progress 1 level (7 levels in all.) Children were seen two times per week. The number of sessions per week is also important. If a child is seen two times per week vs one time, it allows the clinician to move a child forward if a skill is achieved or if the child is struggling it allows the clinician to make adjustments sooner.

It is essential for pediatric voice clinicians to know the cognitive underpinnings of how children learn in order to

successfully choose appropriate vocabulary and tasks which will facilitate successful healthy voice skill acquisition.



References

DeThorne, L. S., Johnson, C. J., Walder, L., & Mahurin-Smith, J. (2009). When "Simon Says" doesn't work: Alternatives to imitation for facilitating early speech development. *American Journal of Speech-Language Pathology*, 18(2), 133-145.

Jacoby, G. P., Lee, L., Kummer, A. W., Levin, L., & Craghead, N. A. (2002). The number of individual treatment units necessary to facilitate functional communication improvements in the speech and language of young children. *American Journal of Speech-Language Pathology*, 11(4), 370-380.

Sawada, M., MOM, S., & Ishii, M. (2002). Effect of metaphorical verbal instruction on modeling of sequential dance skills by young children. *Perceptual and motor skills*, 95(3f), 1097-1105.

Weiss, M. R., Ebbeck, V., & Wiese-Bjornstal, D. M. (1993). Developmental and Psychological Factors Related to Children's Observational Learning of Physical Skills. *Pediatric Exercise Science*, 5(4).

Leslie S. Kessler, MA, CCC-SLP is owner and clinical director of Language & Voice Experience Clinics based in the D.C. region. She received her BA and MA in Speech-Language Pathology and Audiology from New York University. She has advanced training in clinical voice. She has a strong foundation in the performing arts as a trained actor and public speaking trainer. Leslie also published a chapter on pediatric voice therapy, in collaboration with Dr. Katherine Verdolini Abbott.

Leslie has developed proprietary, research-based therapy programs in a variety of domains within speech-language pathology. Included in these interventions is Buzzy Child-Resonant Voice Therapy That's Fun! Buzzy Child has been researched road-tested refined in our clinical practice at Language & Voice Experience, over the past 13 years.

POSTER SESSION SPONSORED BY PENTAX MEDICAL





STROBOSCOPY OSCARS

with your hosts

Michael M. Johns, MD
and Eva van Leer, PhD

Category Winners of the

GOLDEN LARYNX

Patricia Campbell, BS, MM

Patricia Doyle, MA, CCC-SLP

Harry Hollein, PhD

Krzysztof Izdebski, PhD, CCC-SLP

Adam Rubin, MD

Nathan Welham, PhD, CCC-SLP





PRESENTATION OUTREACH COMMITTEE

L-R Front: Brian Gill, Lucille Rubin, Donna Snow, care, Brock Meredith
L-R Back: Deanna Broom, Maria Russo, Elaine Pasqua, Jan Potter Reed, Diane Gaary, Jeannette LoVetri



HEIGH HO! HEIGH HO! TO AVA WORKSHOPS WE GO!



**CHAPTER
AWARD**
South California Chapter,
Lindsay Reder, MD,
President



QUINTANA AWARD

Committee Members Jan Svec, PhD, Mara Behlau, PhD, and Ronald Scherer PhD, with awardee RJ Baken, PhD.

GISELE OLIVERIA AWARD

Co-author
Nico Paulo Paolillo, MD
and Luca Carrozza,
awardee.



Dr. Robert T. Sataloff,
TVF Chairman, presented
Awards at the 2016
Annual Symposium



BEST INTERNATIONAL PRESENTATION

Meike Brockman-Bauser, PhD,
Switzerland



VAN L. LAWRENCE FELLOWSHIP

Matthew Hoch, DMA,
Presented by TVF Chairman,
Dr. Robert Sataloff & representing the
NATS FOUNDATION, NATS President,
Norman Spivey



JOV BEST PAPER AWARD MEDICAL—2015

Miichael Benninger, MD



THE VOICE FOUNDATION

Advancing understanding of the voice through interdisciplinary scientific research & education

THE VOICE FOUNDATION BOARD OF DIRECTORS

CHAIRMAN

Robert Thayer Sataloff, MD,
DMA, FACS

PRESIDENT

Stuart Orsher, MD

VICE-PRESIDENT

Michael S. Benninger, MD

SECRETARY

Mary Hawkshaw

TREASURER AND GENERAL COUNSEL

Brian P. Flaherty, Esq.

ADVISORY BOARD CHAIRMAN

Michael M. Johns III, MD

DIRECTORS

Martina Arroyo
Harolyn Blackwell
Claudia Catania
Jennifer Creed
Abdul-Latif Hamdan, MD
Walt Hunter
Gwen S. Korovin, MD
Justice Sandra Newman
Renata Scotto
Michael Sheehan
George Shirley
Caren J. Sokolow
Diana Soviero

THE VOICE FOUNDATION STAFF

EXECUTIVE DIRECTOR,
THE VOICE FOUNDATION
Maria Russo, MM

EXECUTIVE COORDINATOR
MANAGING EDITOR,
JOURNAL OF VOICE
Katie Erikson, MM



WELCOME NEW BOARD MEMBER

Walt Hunter is a Philadelphia native and an investigative reporter with KYW-TV. He joined the station in 1980 as a crime reporter and was named best television reporter by Philadelphia Magazine in 2004. He has received eight Philadelphia Emmy Awards. The Broadcast Pioneers of Philadelphia inducted Hunter into their Hall of Fame in 2007.

THE VOICE FOUNDATION

219 N. Broad St. 10FL
Philadelphia, PA 19107

(215) 735-7999

(215) 762-5572

voicefoundation.org

office@voicefoundation.org

journal@voicefoundation.org

abstracts@voicefoundation.org

symposium@voicefoundation.org



PENTAX MEDICAL

*World Leaders in Speech,
Voice, and Swallowing
Instrumentation*



**PENTAX
MEDICAL**
Excellence in Focus

www.kaypentax.com

©2013 PENTAX America, Inc. All rights reserved. All company and product names and marks contained within are federally registered trademarks, trademarks or service marks of PENTAX of America, Inc.

MK-442 Flex: A